



Please complete ALL information below and fax your request to 1-888-671-5285

Repatha® Prior Authorization Request Form (Page 1 of 3)

DO NOT COPY FOR FUTURE USE. FORMS ARE UPDATED FREQUENTLY AND MAY BE BARCODED

Member Information (required)			Provider Information (required)		
Member Name:			Provider Name:		
Insurance ID#:			NPI#:	Specialty:	
Date of Birth:			Office Phone:		
Street Address:			Office Fax:		
City:	State:	Zip:	Office Street Address:		
Phone:			City:	State:	Zip:

Medication Information (required)

Medication Name:	Strength:	Dosage Form:
<input type="checkbox"/> Check if generic substitution is acceptable	Directions for Use:	
<input type="checkbox"/> Check if request is for continuation of therapy		

Clinical Information (required)

Select the diagnosis below:

- Homozygous familial hypercholesterolemia (HoFH)
- Primary hyperlipidemia
 - Atherosclerotic cardiovascular disease (ASCVD)
 - Heterozygous familial hypercholesterolemia (HeFH)
 - Secondary prevention of cardiovascular events in patients with ASCVD
- Other diagnosis: _____ ICD-10 Code(s): _____

Clinical Information:

Select if Repatha is prescribed by one of the following specialists:

- Cardiologist
- Endocrinologist
- Lipid specialist

Does the prescriber attest that the information provided is true and accurate to the best of their knowledge and they understand that OptumRx may perform a routine audit and request the medical information necessary to verify the accuracy of the information provided? Yes No

Primary Hyperlipidemia (ASCVD, HeFH, and/or Cardiovascular Prevention):

Select if the patient has heterozygous familial hypercholesterolemia (HeFH) as confirmed by the following:

- Untreated/pre-treatment LDL-cholesterol (LDL-C) > 190 mg/dL in an adult or > 155 mg/dL in a child less than 16 years of age
- Family history of myocardial infarction in first-degree relative less than 60 years of age
- Family history of myocardial infarction in second-degree relative less than 50 years of age
- Family history of LDL-C > 190 mg/dL in first- or second-degree relative
- Family history of familial hypercholesterolemia in first- or second-degree relative
- Family history of tendinous xanthomata and/or arcus cornealis in first- or second-degree relative

Select if there is submission of medical records (e.g., chart notes, laboratory values) documenting the following:*

- Functional mutation in the LDL receptor, ApoB, or PCSK9 gene
- Tendinous xanthomata
- Arcus cornealis before age 45

*Please note: Chart documentation of the above is required to be submitted along with this fax.

<continued on the next page>

This document and others if attached contain information that is privileged, confidential and/or may contain protected health information (PHI). The Provider named above is required to safeguard PHI by applicable law. The information in this document is for the sole use of FutureScripts. Proper consent to disclose PHI between these parties has been obtained. If you received this document by mistake, please know that sharing, copying, distributing or using information in this document is against the law. **If you are not the intended recipient, please notify the sender immediately.**

Repatha® Prior Authorization Request Form (Page 2 of 3)
DO NOT COPY FOR FUTURE USE. FORMS ARE UPDATED FREQUENTLY AND MAY BE BARCODED

<continued from the previous page>

Select if the patient has atherosclerotic cardiovascular disease (ASCVD) confirmed by the following:

- | | |
|---|---|
| <input type="checkbox"/> Acute coronary syndromes | <input type="checkbox"/> Coronary or arterial revascularization |
| <input type="checkbox"/> History of myocardial infarction | <input type="checkbox"/> Stroke |
| <input type="checkbox"/> Stable or unstable angina | <input type="checkbox"/> Transient ischemic attack |
| <input type="checkbox"/> Peripheral arterial disease presumed to be of atherosclerotic origin | |

Has the patient been receiving at least 12 consecutive weeks of **high-intensity** statin therapy [i.e., atorvastatin 40-80 mg, Crestor (rosuvastatin) 20-40 mg]? **Yes** **No**Will the patient continue to receive a **high-intensity** statin at maximally tolerated dose? **Yes** **No**Select if the patient is unable to tolerate **high-intensity** statin therapy as evidenced by one of the following intolerable and persistent (i.e., more than 2 weeks) symptoms:

- Myalgia (muscle symptoms without CK elevations)
- Myositis (muscle symptoms with CK elevations < 10 times upper limit of normal [ULN])

Has the patient been receiving at least 12 consecutive weeks of **moderate-intensity** statin therapy [i.e., atorvastatin 10-20 mg, Crestor (rosuvastatin) 5-10 mg, simvastatin 20-40 mg, pravastatin 40-80 mg, lovastatin 40 mg, Lescol XL (fluvastatin XL) 80 mg, fluvastatin 40 mg twice daily, or Livalo (pitavastatin) 2-4 mg]? **Yes** **No**Will the patient continue to receive a **moderate-intensity** statin at maximally tolerated dose? **Yes** **No**Has the patient been receiving at least 12 consecutive weeks of **low-intensity** statin therapy [i.e., simvastatin 10 mg, pravastatin 10-20mg, lovastatin 20mg, fluvastatin 20-40mg, Livalo (pitavastatin) 1 mg]? **Yes** **No**Will the patient continue to receive a **low-intensity** statin at maximally tolerated dose? **Yes** **No**Select if the patient is unable to tolerate **low-, moderate-, and high-intensity** statin therapy as evidenced by one of the following intolerable and persistent (i.e., more than 2 weeks) symptoms for low-, moderate-, and high-intensity statins:

- Myalgia (muscle symptoms without CK elevations)
- Myositis (muscle symptoms with CK elevations < 10 times upper limit of normal [ULN])

Does the patient have a labeled contraindication to all statins as documented in medical records? **Yes** **No**Has the patient experienced rhabdomyolysis or muscle symptoms with statin treatment with CK elevations > 10 times upper limit of normal? **Yes** **No**Select if there is submission of medical records (e.g., laboratory values) documenting the following LDL-C values while on maximally tolerated lipid-lowering therapy **within the last 120 days**.*

- LDL-C between 70 mg/dL and 99 mg/dL **with** ASCVD
- LDL-C ≥100 mg/dL **with** ASCVD
- LDL-C between 100 mg/dL and 129 mg/dL **without** ASCVD
- LDL-C ≥130 mg/dL **without** ASCVD

Please note: Chart documentation of the above is required to be submitted along with this fax.*Has the patient been receiving at least 12 consecutive weeks of ezetimibe (Zetia) therapy as adjunct to maximally tolerated statin therapy? **Yes **No**If "no" to the above, does the patient have a history of **contraindication or intolerance** to ezetimibe? **Yes** **No****Homozygous Familial Hypercholesterolemia (HoFH):**

Select if there is submission of medical records (e.g., chart notes, laboratory values) documenting the patient has homozygous familial hypercholesterolemia (HoFH) as confirmed by the following.*

- Genetic confirmation of 2 mutations in the LDL receptor, ApoB, PCSK9, or LDL receptor adaptor protein 1 (i.e., LDLRAP1 or ARH)
- Untreated/pre-treatment LDL-cholesterol (LDL-C) > 500 mg/dL
- Treated LDL-C > 300 mg/dL
- Xanthoma before 10 years of age
- Evidence of heterozygous familial hypercholesterolemia in both parents

Please note: Chart documentation of the above is required to be submitted along with this fax.*Is the patient receiving other lipid-lowering therapy (e.g., statin, ezetimibe, LDL, apheresis)? **Yes **No**Will Repatha be used in combination with Juxtapid (lomitapide)? **Yes** **No**



Repatha® Prior Authorization Request Form (Page 3 of 3)
DO NOT COPY FOR FUTURE USE. FORMS ARE UPDATED FREQUENTLY AND MAY BE BARCODED

Reauthorization:

Select if Repatha is prescribed by one of the following specialists:

- Cardiologist
- Endocrinologist
- Lipid specialist

Will medical records (e.g., chart notes, laboratory values) be submitted documenting the patient has had a reduction in LDL-C levels while on PCSK9 therapy?* Yes No

**Please note: Chart documentation of the above is required to be submitted along with this fax.*

Does the prescriber attest that the information provided is true and accurate to the best of their knowledge and they understand that OptumRx may perform a routine audit and request the medical information necessary to verify the accuracy of the information provided? Yes No

For primary hyperlipidemia (ASCVD, HeFH, and/or Cardiovascular Prevention), also answer the following:

Does the patient continue to receive other lipid-lowering therapy (e.g., statins, ezetimibe) at the maximally tolerated dose, unless the patient has documented inability to take these medications? Yes No

For homozygous familial hypercholesterolemia (HoFH), also answer the following:

Does the patient continue to receive other lipid-lowering therapy (e.g., statin, ezetimibe, LDL, apheresis)? Yes No

Will Repatha be used in combination with Juxtapid (lomitapide)? Yes No

Quantity Limit Requests:

What is the quantity requested per MONTH? _____

What is the reason for exceeding the plan limitations?

- Titration or loading dose purposes
- Patient is on a dose-alternating schedule (e.g., one tablet in the morning and two tablets at night, one to two tablets at bedtime)
- Requested strength/dose is not commercially available
- Other: _____

Are there any other comments, diagnoses, symptoms, medications tried or failed, and/or any other information the physician feels is important to this review?

Please note: This request may be denied unless all required information is received.